INFLUENCE OF THE MAGNETIC FIELD ON A VORTEX OF GALLIUM

<u>D. Brito</u>, P. Cardin, H.C. Nataf and G. Marolleau (Département Terre Atmosphère Océan, École Normale Supérieure, Paris, France)

We built an experimental set-up to study the interaction between one single vertical vortex of liquid gallium, generated by a disk, and a transverse magnetic field. This experiment has been conducted on a rotating table. Thus, dominant forces are Coriolis and Lorentz forces, like in the Earth's core. We measured pressure profiles at the top of the vortex, differences in electrical potentials between some points in the vortex and the magnetic field induced by the flow. In order to understand the velocity flow of the vortex, we introduced a simple two dimensional model which actually predicts well these three last different kinds of measurements. As expected, we retrieve that the Elsasser number (ratio of Lorentz over Coriolis forces) is the critical parameter for the dynamical aspect of the vortex. Under the influence of the magnetic field, the vortex is slowed down but remains bidimensionnal; moreover, the size of the vortex increases with the strength of the magnetic field. The magnetic field induced by the loop of electrical current generated in the cylinder, forms a dipole perpendicular to the imposed field. We reach magnetic Reynolds numbers of 0.1 in our experiments.

Daniel Brito École Normale Supérieure Département Terre Atmosphère Océan 24, rue Lhomond 75231 Paris Cedex 05 tel.: 33 1 44 32 22 05 fax.: 33 1 44 32 22 00 e-mail: brito@geophy.ens.fr

Session NP2 : Vortex Dynamics, Turbulence and Transport Processes in Geophysical Flows.Abstract sent to conveners: Dr. David G. Dritschel, Prof. Dr. William K. Dewar and Dr. George G. Sutyrin.Preference for oral presentation.Applied for Young Scientists' Travel Awards.